

Distance regularity in buildings and structure constants in Hecke algebras

Peter Abramenko James Parkinson* Hendrik Van Maldeghem

March 6, 2017

Abstract

In this paper we define generalised spheres in buildings using the simplicial structure and Weyl distance in the building, and we derive an explicit formula for the cardinality of these spheres. We prove a generalised notion of distance regularity in buildings, and develop a combinatorial formula for the cardinalities of intersections of generalised spheres. Motivated by the classical study of algebras associated to distance regular graphs we investigate the algebras and modules of Hecke operators arising from our generalised distance regularity, and prove isomorphisms between these algebras and more well known parabolic Hecke algebras. We conclude with applications of our main results to non-negativity of structure constants in parabolic Hecke algebras, commutativity of algebras of Hecke operators, double coset combinatorics in groups with BN -pairs, and random walks on the simplices of buildings.

Introduction

Buildings are combinatorial/geometric objects with diverse applications in Lie theory and related fields. Spheres and intersections of spheres in buildings (primarily of spherical and affine types) have been studied due to their connections with combinatorial representation theory [21, 12], harmonic analysis on groups of p -adic type [17, 10, 22], incidence structures related to groups of Lie type [8], random walk theory [23], association schemes [8], and number theory [27, 29].

In this paper we define *generalised spheres* in buildings of arbitrary type using the simplicial structure and Weyl distance in the building. We investigate the combinatorics of these spheres, the algebraic structures related to them, and provide applications of our results, including the non-negativity of structure constants in parabolic Hecke algebras, combinatorics of double coset decompositions in groups with BN -pairs, and limit theorems for random walks on buildings.

A main motivation for this paper is to facilitate the study of Hecke algebras related to buildings beyond the spherical and affine cases. Our methodology stems from the theory of *distance regular graphs*, and so we begin by briefly recalling this theory (see [8]). Let $\Gamma = (V, E)$ be a locally finite graph with graph metric $d(\cdot, \cdot)$, and for each $x \in V$ and $k \in \mathbb{N}$ let $F_k(x) = \{y \in V \mid d(x, y) = k\}$ be the sphere of radius k centred at x . The graph Γ is *distance regular* if the intersection cardinality $|F_k(x) \cap F_\ell(y)|$ depends only on k, ℓ and $d(x, y)$. In other words, there are integers $a_{k,\ell}^m$, called the *intersection numbers* of the graph, such that $a_{k,\ell}^m = |F_k(x) \cap F_\ell(y)|$ whenever $d(x, y) = m$. In particular, the cardinality $|F_k(x)| = a_{k,k}^0$ does not depend on $x \in V$. The infinite distance regular graphs are completely classified [14]. In contrast there is no complete classification of finite distance regular graphs, although many partial results exist, for example there are precisely 13 finite cubic distance regular graphs (see [4]).

*Research supported under the Australian Research Council (ARC) discovery grant DP110103205.

A primary tool in the study of finite distance regular graphs is the *Bose-Mesner algebra* of the graph. As a \mathbb{C} -vector space, this algebra is the linear span \mathcal{A} of the matrices $\{A_k \mid k = 0, 1, \dots, d\}$ where d is the diameter of Γ and A_k is the k -adjacency matrix of the graph (with (x, y) -th entry 1 if $d(x, y) = k$ and 0 otherwise). The distance regular property is equivalent to the fact that the matrices A_k satisfy linear relations

$$A_k A_\ell = \sum_{m=0}^d a_{k,\ell}^m A_m \quad \text{for all } k, \ell \in \{0, 1, \dots, d\}. \quad (0.1)$$

It follows that \mathcal{A} is a \mathbb{C} -algebra, and since $a_{k,\ell}^m = a_{\ell,k}^m$ this algebra is commutative.

Let us now return to the setting of the current paper. Let (W, S) be a Coxeter system, and let X be a locally finite, regular building of type (W, S) with chamber set \mathcal{C} and Weyl distance function $\delta : \mathcal{C} \times \mathcal{C} \rightarrow W$. The building X is a labelled simplicial complex, and so each simplex $A \in X$ has a *type* $\tau(A) \subseteq S$. The *cotype* of $A \in X$ is $S \setminus \tau(A)$, and for each $I \subseteq S$ we write X_I for the set of all cotype I simplices of X . For example, a vertex of X has cotype $S \setminus \{s\}$ for some $s \in S$, and a chamber of X has cotype \emptyset .

The Weyl distance function may be extended to arbitrary simplices of X , and if $A \in X_I$ and $B \in X_J$ we have $\delta(A, B) \in R(I, J)$ where $R(I, J)$ is the transversal of minimal length (W_I, W_J) double coset representatives (here W_I and W_J are standard parabolic subgroups of W). For $A \in X_I$ and $w \in R(I, J)$ the *generalised sphere* of “radius” (J, w) centred at A is by definition

$$F(A, J, w) = \{B \in X_J \mid \delta(A, B) = w\}.$$

If the parabolic subgroup W_I is infinite then the cardinality of the sphere $F(A, J, w)$ is often infinite (even in the case of a thin building; that is, a Coxeter complex), and thus we confine our attention to the case that W_I is finite, in which case I is called a *spherical subset* of S .

The main results of this paper are as follows.

- (1) In Theorem 2.1 we prove a new theorem which determines $|F(A, J, w)|$ in the most general situation. Only very special cases were treated before (see [10, Proposition 2.7], [21, Theorem 5.15], [29, Proposition 3.9]), often with considerable effort. Our approach, making use of projections in buildings, is efficient and yields a much more general result.
- (2) Having deduced the most general formula for the numbers $|F(A, J, w)|$, we turn our attention to the considerably more complicated intersection numbers $|F(A, J, u) \cap F(B, J, v)|$. In order to prove a formula for these numbers we introduce the new idea of *pointed pre-galleries* in the Coxeter complex, a model which is inspired by (although distinct from) the alcove walk model developed by Ram [26] and Parkinson, Ram, Schwer [24] for the study of affine Hecke algebras and affine flag varieties. Our formula for the intersection cardinalities is given in Theorem 3.7, and explicit computations using this formula are given in Examples 3.8 and 3.9.
- (3) In Section 4 we define generalised Hecke operators on buildings and investigate the Hecke algebras $\mathcal{A}(I, I)$ and modules $\mathcal{A}(I, J)$ spanned by these operators. These algebras and modules are the natural analogues of the Bose-Mesner algebra of a distance regular graph. In Theorem 4.9 we prove that our Hecke algebras $\mathcal{A}(I, I)$ are isomorphic to the *parabolic Hecke algebras* $\mathbf{1}_I \mathcal{H} \mathbf{1}_I$ introduced in [11], and that our modules $\mathcal{A}(I, J)$ are both left and right modules for parabolic Hecke algebras. Our combinatorial interpretation of the structure constants in these parabolic Hecke algebras yields new insights, for example the positivity result given in Corollary 5.1.

The interpretation of the structure constants in “non-parabolic” Hecke algebras in terms of buildings was made in [21] (this is the case $I = J = \emptyset$ above), and association scheme theoretic interpretations of the structure constants have been developed by Zieschang [33]. In the spherical case implicit connections between Hecke algebras and buildings via finite groups with BN -pairs are given by Iwahori [15], Matsumoto [19], Curtis, Iwahori, and Kilmoyer [11]. Moreover, Brouwer, Cohen and Neumaier [8, Chapter 10], Brouwer and Cohen [7], and Gomi [13] compute structure constants in some spherical cases. In the present paper our point of view is as general as possible. We consider buildings of arbitrary type, with the only assumptions being local finiteness and a very mild regularity hypothesis (which is automatically satisfied in the 2-spherical case). In this generality such buildings may not admit an “interesting” group action, but of course our results also cover the highly symmetric cases including locally finite affine buildings and buildings arising from Kac-Moody groups over finite fields. Also in these cases our formulae for $|F(A, J, w)|$ and $|F(A, J, u) \cap F(B, J, v)|$, and the interpretation of the parabolic Hecke algebras are new.

Other motivations for our investigations include the applications presented in Section 5. In particular we prove a non-negativity result for the structure constants in parabolic Hecke algebras, and classify the algebras of Hecke operators on regular buildings which are commutative. We also obtain combinatorial formulae for double coset cardinalities in groups with BN -pairs, and outline how our algebras generalise the theory of Gelfand pairs associated to buildings. Finally we prove a local limit theorem for a random walk on the simplices of an \tilde{A}_2 building.

1 Background and definitions

1.1 Coxeter groups and complexes

A *Coxeter system* (W, S) is a group W together with a generating set S with defining relations

$$s^2 = 1 \quad \text{and} \quad (st)^{m_{st}} = 1 \quad \text{for all } s, t \in S \text{ with } s \neq t,$$

where $m_{st} = m_{ts} \in \mathbb{Z}_{\geq 2} \cup \{\infty\}$ for all $s \neq t$ (if $m_{st} = \infty$ then it is understood that there is no relation between s and t). We shall always assume in this paper that the *rank* of (W, S) , defined as $|S|$, is finite.

The *length* of $w \in W$ is

$$\ell(w) = \min\{n \geq 0 \mid w = s_1 \cdots s_n \text{ with } s_1, \dots, s_n \in S\}.$$

We say that a decomposition $w = s_1 \cdots s_n$ with $s_1, \dots, s_n \in S$ is *reduced* if $n = \ell(w)$. The *Bruhat order* on W can be defined as follows: For $u, w \in W$, $u < w$ if and only if a reduced decomposition of u can be obtained by deleting some of the s_i in a reduced decomposition $w = s_1 \cdots s_n$ of w (so that in particular $\ell(u) < \ell(w)$).

A *standard subgroup* of W is by definition a subgroup of the form $W_I = \langle \{s \mid s \in I\} \rangle$ with I a subset of S . Such a subset I is called *spherical* if W_I is finite. To every Coxeter system (W, S) one associates the *Coxeter complex* $\Sigma(W, S)$, see [2, Section 3.1]. Thus $\Sigma(W, S)$ is a thin chamber complex, its chambers are identified with the elements of W , its simplices with standard cosets wW_I , and wW_I is a face of uW_J if and only if $uW_J \subseteq wW_I$ (for $u, w \in W$ and $I, J \subseteq S$). The *type* of a simplex wW_I in $\Sigma(W, S)$ is by definition $S \setminus I$ and its *cotype* is I .

1.2 Double cosets and reduced elements

Let (W, S) be a Coxeter system. The following facts about cosets and double cosets in W are well-known, see for instance [2, Subsection 2.3.2]:

- (a) If $J \subseteq S$ and $w \in W$ then the coset wW_J has a unique minimal length representative w_1 , and $\ell(w_1y) = \ell(w_1) + \ell(y)$ for all $y \in W_J$.
- (b) If $I, J \subseteq S$ and $w \in W$ then the double coset W_IwW_J has a unique minimal length representative. This representative is called (I, J) -reduced, and we let

$$R(I, J) = \{w \in W \mid w \text{ is } (I, J)\text{-reduced}\}.$$

Note that $R(I, J)$ indexes the decomposition of W into W_IwW_J double cosets.

- (c) In general it is not true that $\ell(xwy) = \ell(x) + \ell(w) + \ell(y)$ for all $x \in W_I, y \in W_J$ and $w \in R(I, J)$. In order to address this issue, we introduce the following notation, where I and J are subsets of S and $w \in R(I, J)$:

$$W_I(J, w) = W_I \cap wW_Jw^{-1}$$

$M_I(J, w)$ = the set of all minimal length representatives of cosets in $W_I/W_I(J, w)$.

Note that $W_I(J, w) = \{v \in W_I \mid vwW_J = wW_J\}$ is the subgroup of W_I stabilising wW_J , and by [2, Lemma 2.25] we have that

$$W_I(J, w) = W_{I \cap wJw^{-1}}. \tag{1.1}$$

Thus applying point (a) it follows that each $v \in W_I$ can be uniquely expressed in the form

$$v = xz \text{ with } x \in M_I(J, w), z \in W_I(J, w), \text{ and that } \ell(v) = \ell(xz) = \ell(x) + \ell(z). \tag{1.2}$$

We now obtain an analogous result for double cosets.

Lemma 1.1. *Let I and J be subsets of S and let $w \in R(I, J)$. Then each $v \in W_IwW_J$ can be written in exactly one way as*

$$v = xwz \quad \text{with } x \in M_I(J, w) \quad \text{and } z \in W_J.$$

Moreover, if $x \in M_I(J, w)$, $w \in R(I, J)$, and $z \in W_J$ then $\ell(xwz) = \ell(x) + \ell(w) + \ell(z)$.

Proof. It follows immediately from $W_I = M_I(J, w)W_I(J, w)$ and $W_I(J, w) = W_I \cap wW_Jw^{-1}$ that $W_IwW_J = M_I(J, w)wW_J$. If $x, x' \in M_I(J, w)$ and $y, y' \in W_J$, then $xwy = x'wy'$ implies $x^{-1}x' = wyy'^{-1}w^{-1} \in W_I(J, w)$, and so $x' = x$ is the unique minimal length representative of $xW_I(J, w)$, which then also implies $y' = y$. Finally, assume that $\ell(xwz) < \ell(x) + \ell(w) + \ell(z)$. Using the deletion condition, we find $x' \in W_I, y' \in W_J$ with $x' < x, y' < y$ and $x'wy' = xwy$ (see [2, Lemma 2.24]). However, as before $xwy = x'wy'$ implies $xW_I(J, w) = x'W_I(J, w)$, contradicting the minimality of x in $xW_I(J, w)$. \square

1.3 Buildings

We assume that the reader is already acquainted with the theory of buildings, and our main reference for this subject is [2, Chapters 4 and 5]. In this paper, buildings will mainly be considered as chamber (and hence as simplicial) complexes but the associated Weyl distance will also play an important role.

In the following, X will always denote a building and \mathcal{C} its set of chambers. The apartments of X are all isomorphic to the standard Coxeter complex $\Sigma(W, S)$ for a Coxeter system (W, S) which is determined, up to isomorphism, by X . We then say that X is of *type* (W, S) and

of rank $|S|$. If we fix an apartment \mathcal{A} of X and an isomorphism $f : \Sigma(W, S) \rightarrow \mathcal{A}$, the *type* $\tau(A)$ of a simplex A of \mathcal{A} is by definition the type of $f^{-1}(A)$. There is then a unique extension $\tau : X \rightarrow \{I \mid I \subseteq S\}$ such that $\tau(A) \subseteq \tau(B)$ whenever A is a face of B and $\tau(B) = S$ if and only if B is a chamber of X . We fix such a type function on X and define the *cotype* of a simplex $A \in X$ to be $S \setminus \tau(A)$. We set

$$X_I = \{A \in X \mid A \text{ is of cotype } I\}.$$

For $c \in \mathcal{C}$ and $I \subseteq S$ the unique element of X_I which is a face of c is denoted by c_I . For $A \in X$ let

$$\mathcal{C}(A) = \{c \in \mathcal{C} \mid A \text{ is a face of } c\},$$

and for any subset $M \subset X$ we write $\mathcal{C}(M) = \bigcup_{A \in M} \mathcal{C}(A)$. If $A \in X_I$ then $\mathcal{C}(A) = \{c \in \mathcal{C} \mid c_I = A\}$. *Panels* of X are codimension 1 simplices; an *s-panel* is one of cotype $\{s\}$ for $s \in S$. The building X is called *thick* if each panel is contained in at least 3 chambers.

A *pregallery* in X of length $n \in \mathbb{N}_0$ is a sequence

$$\gamma = (c_0, s_1, c_1, s_2, c_2, s_3, \dots, s_n, c_n)$$

where $s_1, \dots, s_n \in S$ and $c_0, c_1, \dots, c_n \in \mathcal{C}$ and, for each $j = 1, \dots, n$, either $c_{j-1} = c_j$ or $c_{j-1} \cap c_j$ is an s_j -panel. The *type* of the pregallery $\gamma = (c_0, s_1, c_1, s_2, \dots, s_n, c_n)$ is (s_1, \dots, s_n) . The type is called *reduced* if $w = s_1 \cdots s_n \in W$ satisfies $\ell(w) = n$. The chamber c_0 is called the *start* of γ , and the chamber c_n is called the *end* of γ , written $\text{end}(\gamma) = c_n$.

A *gallery* of type (s_1, \dots, s_n) is a pregallery $\gamma = (c_0, s_1, c_1, s_2, \dots, s_n, c_n)$ such that $c_{j-1} \neq c_j$ for all $j = 1, \dots, n$. Note that the type of the gallery γ is already determined by the sequence (c_0, c_1, \dots, c_n) , although this is not true for pregalleries. For $c, d \in \mathcal{C}$ the *gallery distance* $\text{dist}(c, d)$ is the minimal length of galleries starting in c and ending in d .

For any two chambers $c, d \in \mathcal{C}$ there exists an element $w \in W$ with the following property: There exists a gallery γ of reduced type (s_1, \dots, s_n) which starts in c and ends in d if and only if $s_1 \cdots s_n = w$. We set $\delta(c, d) = w$ for this element w , thus obtaining a *Weyl distance* function $\delta : \mathcal{C} \times \mathcal{C} \rightarrow W$. For the basic properties of δ we refer to [2, Section 4.8]. We mention in passing that $\text{dist}(c, d) = \ell(\delta(c, d))$. For $A \in X_I$ and $B \in X_J$ (with $I, J \subseteq S$) we have

$$\delta(\mathcal{C}(A) \times \mathcal{C}(B)) = W_I w W_J$$

whenever $w = \delta(c, d)$ for fixed chambers $c \in \mathcal{C}(A)$ and $d \in \mathcal{C}(B)$. This allows us to extend the Weyl distance to arbitrary simplices of X by setting (for $A \in X_I$ and $B \in X_J$):

$$\delta(A, B) = \text{the unique } (I, J)\text{-reduced element of } \delta(\mathcal{C}(A) \times \mathcal{C}(B)). \quad (1.3)$$

1.4 Retractions and projections

We now briefly recall two fundamental concepts of building theory which we will apply later. First, for a fixed apartment \mathcal{A} of X and a fixed chamber a of \mathcal{A} , there is the *retraction* $\rho_{\mathcal{A}, a}$ of X onto \mathcal{A} centred at a . It is the type-preserving simplicial map from X to \mathcal{A} with the property that for every apartment \mathcal{A}' containing a the restriction of $\rho_{\mathcal{A}, a}$ to \mathcal{A}' is the unique type-preserving isomorphism of \mathcal{A}' onto \mathcal{A} fixing a . From this one readily deduces the following facts:

(R1) For all $c \in \mathcal{C}$ we have $\delta(a, c) = \delta(a, \rho_{\mathcal{A}, a}(c))$.

(R2) If $\gamma = (c_0, \dots, c_n)$ is a gallery of type (s_1, \dots, s_n) in X , then the image

$$\rho_{\mathcal{A},a}(\gamma) = (\rho_{\mathcal{A},a}(c_0), s_1, \dots, s_n, \rho_{\mathcal{A},a}(c_n))$$

is a pregallery of the same type in \mathcal{A} .

We now turn to projections. All of the properties relevant for us can be found in [2, Section 4.9]. For any simplex $A \in X$ and any chamber $c \in \mathcal{C}$, there is a unique chamber in $\mathcal{C}(A)$, called the *projection of c onto A* and denoted by $\text{proj}_A(c)$, such that $\text{dist}(\text{proj}_A(c), c) < \text{dist}(d, c)$ for all $d \in \mathcal{C}(A) \setminus \{\text{proj}_A(c)\}$. For any two simplices $A, B \in X$, there is a unique simplex containing A , called the *projection $\text{proj}_A(B)$ of B onto A* , such that $\{\text{proj}_A(c) \mid c \in \mathcal{C}(B)\} = \mathcal{C}(\text{proj}_A(B))$. We shall use the following properties of projections:

(P1) If $A \in X_I, B \in X_J$ and $w = \delta(A, B)$, then $\text{proj}_A(B)$ has cotype $I \cap wJw^{-1}$. Furthermore,

$$\{c \in \mathcal{C}(A) \mid \text{there exists } d \in \mathcal{C}(B) \text{ such that } \delta(c, d) = w\} = \mathcal{C}(\text{proj}_A(B)).$$

(P2) There exist bijections (induced by projection maps) between $\mathcal{C}(\text{proj}_A(B))$ and $\mathcal{C}(\text{proj}_B(A))$.

1.5 Generalised spheres in regular buildings

Spheres (and balls) in buildings can be defined using the gallery distance. In this paper we study the refinement of this notion by considering spheres defined using the Weyl distance. For a building X of type (W, S) , $c \in \mathcal{C}$ and $w \in W$, let

$$\mathcal{C}_w(c) = \{d \in \mathcal{C} \mid \delta(c, d) = w\}.$$

And for $I, J \subseteq S, A \in X_I$ and $w \in R(I, J)$, we define

$$F(A, J, w) = \{B \in X_J \mid \delta(A, B) = w\}.$$

That is what we mean by a *generalised sphere*. Note that for $c \in \mathcal{C}$ and $w \in W = R(\emptyset, \emptyset)$ we have $F(c, \emptyset, w) = \mathcal{C}_w(c)$. It is one of the purposes of the present paper to compute the cardinalities $|F(A, J, w)|$ for regular, locally finite buildings and spherical cotype I (if I is not spherical then the cardinality $|F(A, J, w)|$ is often infinite).

A building X is called *regular* if there is a bijection between $\mathcal{C}(\pi)$ and $\mathcal{C}(\pi')$ whenever π and π' are panels of the same cotype. If X is thick and $m_{st} < \infty$ for all $s, t \in S$ then X is necessarily regular (see for example, [21, Theorem 2.4]). If each panel is contained in only finitely many chambers then we call X *locally finite*. For the remainder of this paper we assume that X is a regular, locally finite building of type (W, S) .

For each $s \in S$ we set

$$q_s + 1 = |\mathcal{C}(\pi_s)| \quad \text{for any } s\text{-panel } \pi_s.$$

The *parameters* of X are the integers $(q_s)_{s \in S}$. Note that $q_s = |\mathcal{C}_s(c)|$ for any $c \in \mathcal{C}$. Induction on $\ell(w)$ shows that if $w \in W$ has a reduced decomposition $w = s_1 \cdots s_\ell$, then

$$|\mathcal{C}_w(c)| = q_{s_1} \cdots q_{s_\ell}, \tag{1.4}$$

see [21, Proposition 2.1]. In particular, the number $q_w = q_{s_1} \cdots q_{s_\ell} = |\mathcal{C}_w(c)|$ (with the convention that $q_1 = 1$) does not depend on the choice of the chamber $c \in \mathcal{C}$ or the reduced expression

for w . Therefore $q_{uv} = q_u q_v$ whenever $\ell(uv) = \ell(u) + \ell(v)$. It also follows that $q_s = q_t$ whenever the order of st in W is finite and odd (but we will not explicitly use this here).

If $A \in X_I$ and $c \in \mathcal{C}(A)$ then

$$\mathcal{C}(A) = \bigsqcup_{w \in W_I} \mathcal{C}_w(c).$$

Thus if I is spherical and $A \in X_I$ then the cardinality $N(I) := |\mathcal{C}(A)|$ is finite and does not depend on $A \in X_I$, since

$$N(I) = |\mathcal{C}(A)| = \sum_{w \in W_I} q_w. \quad (1.5)$$

If $I, J \subseteq S$ with I spherical, $A \in X_I$, $B \in X_J$ and $w = \delta(A, B)$ then properties (P1) and (P2) in Section 1.4 imply that

$$N(I \cap wJw^{-1}) = N(J \cap w^{-1}Iw). \quad (1.6)$$

2 Counting simplices at given Weyl distance

In the following theorem we give our explicit formula for the cardinality of generalised spheres in a regular building.

Theorem 2.1. *Let I be a spherical subset of S , J a subset of S , and suppose that $A \in X_I$. If $w \in R(I, J)$ then*

$$|F(A, J, w)| = \frac{N(I)}{N(I \cap wJw^{-1})} q_w,$$

where $N(K)$, for K spherical, is given by (1.5).

Proof. Let $c \in \mathcal{C}(A)$ and $d \in \mathcal{C}_w(c)$. Then $d_J \in F(A, J, w)$. If $c' \in \mathcal{C}(A)$ with $c' \neq c$ then $\mathcal{C}_w(c) \cap \mathcal{C}_w(c') = \emptyset$, for if $d \in \mathcal{C}_w(c)$ and $\delta(c', c) = w_I \in W_I$ then $\delta(c', d) = w_I w$ (since $w \in R(I, J)$). Hence if $d \in \mathcal{C}_w(c)$ then $d \notin \mathcal{C}_w(c')$.

The above observations show that there is a well defined map

$$f : \bigsqcup_{c \in \mathcal{C}(A)} \mathcal{C}_w(c) \rightarrow F(A, J, w), \quad \text{given by } f(d) = d_J.$$

By (1.4) and (1.5) the domain of f has cardinality $N(I)q_w$. We now compute the cardinalities of the fibres of f . For $B \in F(A, J, w)$ we have

$$f^{-1}(B) = \{d \in \mathcal{C}(B) \mid \text{there exists } c \in \mathcal{C}(A) \text{ such that } \delta(c, d) = w\},$$

and by (P1) in Section 1.4 it follows that

$$f^{-1}(B) = \{d \in \mathcal{C}(B) \mid \text{proj}_B(A) \subseteq d\} = \mathcal{C}(\text{proj}_B(A)).$$

The cotype of the simplex $\text{proj}_B(A)$ is $J \cap w^{-1}Iw$ (see (P1) in Section 1.4), and thus by (1.5) we have $|f^{-1}(B)| = N(J \cap w^{-1}Iw) \geq 1$. This shows firstly that f is surjective, and secondly that

$$|F(A, J, w)| = \frac{N(I)q_w}{N(J \cap w^{-1}Iw)}.$$

The claim now follows since $N(J \cap w^{-1}Iw) = N(I \cap wJw^{-1})$ by (1.6). \square

Remark 2.2. We make the following remarks.

- (a) In the case of *special vertices* of a fixed type in an affine building, our formula in Theorem 2.1 recovers [10, Proposition 2.7] (in the case of \tilde{A}_n buildings) and [21, Theorem 5.15] (for general affine buildings).
- (b) We outline a representation theoretic proof of Theorem 2.1 in Remark 4.7. An advantage of the building theoretic proof presented above is that a combinatorial significance is attached to the denominator $N(I \cap wJw^{-1})$ in the formula.
- (c) We note that for all finite Coxeter groups W_I there is an explicit closed formula for the polynomial $N(I)$ (this polynomial is often called the *Poincaré polynomial* of W_I). See, for example [18]. Thus the formula in Theorem 2.1 can be made completely explicit.

The *gallery distance* between simplices A and B is $\text{dist}(A, B) = \ell(\delta(A, B))$. The following elementary corollary gives formulae for the cardinalities of spheres with respect to this numerical distance.

Corollary 2.3. *If I is a spherical subset of S , J is a subset of S , and $A \in X_I$, then*

$$|\{B \in X_J \mid \text{dist}(A, B) = n\}| = \sum_{\{w \in R(I, J) \mid \ell(w) = n\}} \frac{N(I)}{N(I \cap wJw^{-1})} q_w. \quad (2.1)$$

In particular, if $A \in X_I$ is a vertex of spherical cotype $I = S \setminus \{s\}$, and if B is a vertex of X with $\text{dist}(A, B) = 1$, then B is a vertex of cotype I and

$$|\{B \in X_I \mid \text{dist}(A, B) = 1\}| = \frac{N(I)q_s}{N(I \cap sIs)} = \frac{N(I)q_s}{N(\{t \in I \mid st = ts\})}. \quad (2.2)$$

Proof. Equation (2.1) follows immediately from Theorem 2.1. For equation (2.2), note that if $A \in X_I$ is a vertex of cotype $I = S \setminus \{s\}$ and if $B \in X_J$ is a vertex of cotype $J = S \setminus \{s'\}$ with $\text{dist}(A, B) = 1$ then $\delta(A, B) = t$ for some $t \in S$ with $t \in R(I, J)$. Thus $t \notin I \cup J$, and hence $I = J$ and $t = s$. Thus B is a vertex of cotype I , and if I is spherical then equation (2.1) gives $|\{B \in X_I \mid \text{dist}(A, B) = 1\}| = N(I)q_s/N(I \cap sIs)$. It is elementary that $I \cap sIs = \{t \in I \mid st = ts\}$, hence (2.2). \square

In the literature [27, 29] computing the cardinality of the sphere of vertices at numerical distance 1 from a given vertex has been an issue even for classical affine buildings of types \tilde{A}_n and \tilde{C}_n over local fields. Corollary 2.3, and in particular (2.2), gives a very general and explicit solution to this problem.

Example 2.4. Let X be an affine building of type \tilde{A}_n with thickness q . By (2.2) the number of vertices at numerical distance 1 from a given vertex of cotype $I = S \setminus \{s\}$ is

$$\frac{N(I)}{N(I \cap sIs)} q = \frac{|A_n(q)|}{|A_{n-2}(q)|} q = \frac{(q^{n+1} - 1)(q^n - 1)}{(q - 1)^2} q, \quad (2.3)$$

where we use the elementary formula for the number of chambers of an A_n building $A_n(q)$ with thickness q (or, alternatively, the well known closed form for the Poincaré polynomial). The formula (2.3) was conjectured by Schwartz and Shemanske in [27], and proven by Setyadi in [29].

Example 2.5. Let X be an affine building of type \tilde{C}_n , and assume that $q_s = q$ for all $s \in S$.

(a) Suppose first that $s \in S$ is a special type (that is, s is an end node of the Coxeter diagram). Then by (2.2) the number of vertices at numerical distance 1 from a given cotype $I = S \setminus \{s\}$ vertex is

$$\frac{N(I)}{N(I \cap sIs)}q = \frac{|C_n(q)|}{|C_{n-1}(q)|}q = \frac{q^{2n} - 1}{q - 1}q,$$

where we use elementary and well known formulae for the number of chambers of a $C_n(q)$ building with thickness q . This recovers [29, Proposition 3.9].

(b) Suppose now that $s \in S$ is not a special type. If s is at length k from an end of the Coxeter diagram, then $W_I = C_k \times C_{n-k}$ and $W_{I \cap sIs} = C_{k-1} \times C_{n-k-1}$, and hence the number of vertices at numerical distance 1 from a given vertex of cotype $I = S \setminus \{s\}$ is

$$\frac{N(I)}{N(I \cap sIs)}q = \frac{|C_k(q)| \times |C_{n-k}(q)|}{|C_{k-1}(q)| \times |C_{n-k-1}(q)|}q = \frac{(q^{2k} - 1)(q^{2(n-k)} - 1)}{(q - 1)^2}q.$$

3 Intersection numbers in regular buildings

Let $\Gamma = (V, E)$ be a locally finite graph, and for $k \in \mathbb{N}$ let $F_k(x) = \{y \in V \mid d(x, y) = k\}$ be the sphere of radius k centred at $x \in V$. As recalled in the introduction, the graph Γ is *distance regular* if for all $k, \ell \in \mathbb{N}$ we have $|F_k(x) \cap F_\ell(y)| = |F_k(x') \cap F_\ell(y')|$ whenever $d(x, y) = d(x', y')$. In other words, there are numbers $a_{k,\ell}^m \in \mathbb{N}$ such that

$$a_{k,\ell}^m = |F_k(x) \cap F_\ell(y)| \quad \text{whenever } y \in F_m(x).$$

Distance regular graphs have been extensively studied, and have a rich algebraic theory centring around the *intersection numbers* $a_{k,\ell}^m$ (see for example [8]).

In this section we prove a generalised notion of distance regularity for regular buildings using the Weyl distance between simplices. We also develop a combinatorial formula for the corresponding intersection numbers. Consider first the simplest case, where the simplices are chambers. By [21, Proposition 3.9] there are numbers $c_{u,v}^w \in \mathbb{N}$ ($u, v, w \in W$) such that

$$c_{u,v}^w = |\mathcal{C}_u(a) \cap \mathcal{C}_v(b)| \quad \text{whenever } a, b \in \mathcal{C} \text{ with } b \in \mathcal{C}_w(c). \quad (3.1)$$

We now extend this chamber distance regularity to a general distance regularity result for regular buildings.

Theorem 3.1. *Let I, J, K be spherical subsets of S , and suppose that $u \in R(I, J)$, $v \in R(K, J)$, and $w \in R(I, K)$. There are numbers $c_{u,v}^w(I, J, K) \in \mathbb{N}$ such that*

$$c_{u,v}^w(I, J, K) = |F(A, J, u) \cap F(B, J, v)| \quad \text{whenever } A \in X_I \text{ and } B \in F(A, K, w).$$

Moreover, for any $z \in W_I W_K$ we have

$$c_{u,v}^w(I, J, K) = \frac{1}{N(J)} \sum_{\substack{x \in W_I u W_J \\ y \in W_K v W_J}} c_{x,y}^z. \quad (3.2)$$

where the numbers $c_{x,y}^z \in \mathbb{N}$ are from (3.1).

Proof. Let $A \in X_I$ and $B \in F(A, K, w)$, and let a, b be any chambers with $a \in \mathcal{C}(A)$ and $b \in \mathcal{C}(B)$ and set $z = \delta(a, b)$. We claim that

$$\mathcal{C}(F(A, J, u) \cap F(B, J, v)) = \bigsqcup_{\substack{x \in W_I u W_J \\ y \in W_K v W_J}} \mathcal{C}_x(a) \cap \mathcal{C}_y(b).$$

For if $d \in \mathcal{C}(F(A, J, u) \cap F(B, J, v))$ then $D = d_J \in F(A, J, u) \cap F(B, J, v)$ and so $\delta(A, D) = u$ and $\delta(B, D) = v$. Thus $\delta(a, d) \in W_I u W_J$ and $\delta(b, d) \in W_K v W_J$. Conversely, if $d \in \mathcal{C}_x(a) \cap \mathcal{C}_y(b)$ with $x \in W_I u W_J$ and $y \in W_K v W_J$ and $D = d_J$ then $\delta(A, D) = u$ and $\delta(B, D) = v$. Thus $D \in F(A, J, u) \cap F(B, J, v)$, and so $d \in \mathcal{C}(D) \subseteq \mathcal{C}(F(A, J, u) \cap F(B, J, v))$.

Hence, by (3.1), we have

$$|\mathcal{C}(F(A, J, u) \cap F(B, J, v))| = \sum_{\substack{x \in W_I u W_J \\ y \in W_K v W_J}} c_{x,y}^z.$$

Since each J -simplex of $F(A, J, u) \cap F(B, J, v)$ is contained in precisely $N(J)$ chambers of $\mathcal{C}(F(A, J, u) \cap F(B, J, v))$ formula (3.2) holds. Thus $|\mathcal{C}(F(A, J, u) \cap F(B, J, v))|$ depends only on u, v and w (for we could choose $a \in \mathcal{C}(A)$ and $b \in \mathcal{C}(B)$ with $\delta(a, b) = w$). \square

Remark 3.2. If I, J, K are spherical and if $u \in R(I, J)$, $v \in R(K, J)$, and $w \in R(I, K)$ then $c_{u,v}^w(I, J, K) = 0$ if $\ell(w) > \ell(u) + \ell(v) + \ell(w_J)$, where $w_J \in W_J$ is the longest element of W_J . To see this, note that if $A \in X_I$ and $B \in X_K$ with $\delta(A, B) = w$, and if $D \in F(A, J, u) \cap F(B, J, v)$, then there are chambers $a \in \mathcal{C}(A)$, $d_1, d_2 \in \mathcal{C}(D)$, and $b \in \mathcal{C}(B)$ with $\delta(a, d_1) = u$ and $\delta(d_2, b) = v^{-1}$. Thus $\ell(w) = \ell(\delta(A, B)) \leq \ell(u) + \ell(\delta(d_1, d_2)) + \ell(v^{-1}) \leq \ell(u) + \ell(v) + \ell(w_J)$.

The numbers $c_{u,v}^w(I, J, K)$ are called *intersection numbers*. The following refined formula for the intersection numbers will be used for our combinatorial formula in Theorem 3.7. It appears that the formula in Proposition 3.3 does not easily follow from the formula in Theorem 3.1 by obvious algebraic manipulations, despite the apparent similarity in the formulae. Thus we give a direct and independent proof below.

Proposition 3.3. *Let $I, J, K \subseteq S$ be spherical and suppose that $u \in R(I, J)$, $v \in R(K, J)$, and $w \in R(I, K)$. For any $z \in W_I w W_K$ we have (with $M_K(J, v)$ as in Section 1.2):*

$$c_{u,v}^w(I, J, K) = \sum_{\substack{x \in W_I u W_J \\ y \in M_K(J, v)v}} c_{x,y}^z.$$

Proof. Choose $A \in X_I$ and $B \in F(A, K, w)$, and fix chambers $a \in \mathcal{C}(A)$ and $b \in \mathcal{C}(B)$ such that $\delta(a, b) = z$. We claim that the mapping $c \mapsto c_J$ gives a bijection

$$\bigsqcup_{\substack{x \in W_I u W_J \\ y \in M_K(J, v)v}} \mathcal{C}_x(a) \cap \mathcal{C}_y(b) \rightarrow F(A, J, u) \cap F(B, J, v).$$

To check surjectivity, let $D \in F(A, J, u) \cap F(B, J, v)$. Since $b \in \mathcal{C}(B)$ and $\delta(\mathcal{C}(B) \times \mathcal{C}(D)) = W_K v W_J = M_K(v, J) v W_J$ we have $\delta(\{b\} \times \mathcal{C}(D)) = m v W_J$ for some $m \in M_K(J, v)$. Thus there is a chamber $d \in \mathcal{C}(D)$ with $\delta(b, d) = m v \in M_K(J, v)v$. Since $d \in \mathcal{C}(D)$ and D has cotype J , we have $d_J = D$. Since $a \in \mathcal{C}(A)$, $d \in \mathcal{C}(D)$ and $\delta(A, D) = u$ we have $\delta(a, d) \in W_I u W_J$.

To check injectivity, suppose that $d, d' \in \mathcal{C}$ with $\delta(b, d) = m v \in M_K(J, v)v$ and $\delta(b, d') = m' v \in M_K(J, v)v$ and $d_J = d'_J$. Then $\delta(d, d') \in W_J$, and so $m' v = \delta(b, d') \in \delta(b, d) W_J$. Thus $m' v = m v w_J$ for some $w_J \in W_J$, and it follows from Lemma 1.1 that $m' = m$ and $w_J = 1$. Thus $d = d'$. The formula now follows by taking cardinalities and using (3.1). \square

We now develop a combinatorial formula for the intersection cardinalities $c_{u,v}^w(I, J, K)$ in terms of “pointed pregalleries” (see the definition in the following paragraph). These combinatorial objects are inspired by the alcove walk model developed by Ram [26]. For the readers familiar with this theory, our pointed pregalleries are analogues of positively folded alcove walks where instead of the ‘folding’ occurring away from a point at infinity, instead the folding occurs away from a fixed chamber. Note that our pointed pregalleries are defined for all Coxeter systems, whereas positively folded alcove walks are only defined for affine Coxeter systems.

Let \mathcal{A} be an apartment of X and let a be a chamber of \mathcal{A} . An a -pointed pregallery in \mathcal{A} of type (s_1, \dots, s_n) is a pregallery $\gamma = (c_0, s_1, c_1, s_2, \dots, s_n, c_n)$ in \mathcal{A} such that

$$\text{if } c_{j-1} = c_j \text{ then } \delta(a, c_{j-1})s_j < \delta(a, c_j).$$

Our interest in a -pointed pregalleries stems from the following elementary fact.

Lemma 3.4. *Let $a, b \in \mathcal{C}$ and let \mathcal{A} be an apartment with $a, b \in \mathcal{A}$. If Γ is a gallery in X of type \mathbf{v} starting at b then $\rho_{\mathcal{A},a}(\Gamma)$ is an a -pointed pregallery in \mathcal{A} of type \mathbf{v} starting at b*

Proof. Write $\rho = \rho_{\mathcal{A},a}$. In view of (R2) in Section 1.4 it suffices to show that $\rho(\Gamma)$ is a -pointed. Write $\Gamma = (b_0, s_1, \dots, s_n, b_n)$ and suppose that $\rho(b_{j-1}) = \rho(b_j)$. If $\delta(a, \rho(b_{j-1}))s_j > \delta(a, \rho(b_{j-1}))$ then $\delta(a, b_{j-1})s_j > \delta(a, b_{j-1})$ by (R1), and hence $\delta(a, b_j) = \delta(a, b_{j-1})s_j$. Thus by (R1) we have $\delta(a, \rho(b_j)) = \delta(a, \rho(b_{j-1}))s_j$, contradicting $\rho(b_{j-1}) = \rho(b_j)$. Thus $\delta(a, \rho(b_{j-1}))s_j < \delta(a, \rho(b_{j-1}))$ and so $\rho(\Gamma)$ is a -pointed. \square

If $\gamma = (c_0, s_1, \dots, s_n, c_n)$ is an a -pointed pregallery in an apartment \mathcal{A} , and $s \in S$ then we define

$$\begin{aligned} \alpha_s(\gamma) &= |\{j \mid s_j = s \text{ and } \delta(a, c_{j-1}) < \delta(a, c_j)\}| \\ \sigma_s(\gamma) &= |\{j \mid s_j = s \text{ and } c_j = c_{j-1}\}|. \end{aligned}$$

(Intuitively, $\alpha_s(\gamma)$ counts the number of “ascent steps” in the pointed pregallery that occur on cotype s panels, where ascent step means that the gallery increases length from a , and $\sigma_s(\gamma)$ counts the number of stutters in the gallery that occur on cotype s panels).

If γ is an a -pointed pregallery we define

$$q(\gamma) = \prod_{s \in S} q_s^{\alpha_s(\gamma)} (q_s - 1)^{\sigma_s(\gamma)}.$$

Proposition 3.5. *Let X be a thick regular building of type W with parameters $(q_s)_{s \in S}$. Let $a, b \in \mathcal{C}$ with $\delta(a, b) = w$ and let \mathcal{A} be an apartment containing a and b . Let $\mathbf{v} = (s_1, \dots, s_n) \in S^n$ and define*

$$\begin{aligned} \mathcal{G}(b, \mathbf{v}) &= \{\text{galleries in } X \text{ of type } \mathbf{v} \text{ starting at } b\}, \\ \mathcal{P}_{\mathcal{A},a}(b, \mathbf{v}) &= \{a\text{-pointed pregalleries in } \mathcal{A} \text{ of type } \mathbf{v} \text{ starting at } b\}. \end{aligned}$$

Then

1. $\rho_{\mathcal{A},a}(\mathcal{G}(b, \mathbf{v})) = \mathcal{P}_{\mathcal{A},a}(b, \mathbf{v})$, and
2. $|\rho_{\mathcal{A},a}^{-1}(\gamma)| = q(\gamma)$ for all $\gamma \in \mathcal{P}_{\mathcal{A},a}(b, \mathbf{v})$

Proof. Write $\rho = \rho_{\mathcal{A},a}$. From Lemma 3.4 we have $\rho(\mathcal{G}(b, \mathbf{v})) \subseteq \mathcal{P}_{\mathcal{A},a}(b, \mathbf{v})$. We prove the remaining statements by induction on n . Thus suppose that the result is true for $\mathbf{v}' = (s_1, \dots, s_n)$ and let $\mathbf{v} = (s_1, \dots, s_n, t)$. Let

$$\gamma = (c_0, s_1, c_1, \dots, s_n, c_n, t, c_{n+1}) \in \mathcal{P}_{\mathcal{A},a}(b, \mathbf{v})$$

(and so $c_0 = b$) and let $\gamma' = (c_0, s_1, c_1, \dots, s_n, c_n) \in \mathcal{P}_{\mathcal{A},a}(b, \mathbf{v}')$. By the induction hypothesis there are $q(\gamma')$ galleries $\Gamma' = (b_0, s_1, \dots, s_n, b_n) \in \mathcal{G}(b, \mathbf{v}')$ with $\rho(\Gamma') = \gamma'$ (that is, $\rho(b_j) = c_j$ for all $0 \leq j \leq n$). For each such gallery Γ' the number of ways of extending Γ' to a gallery $\Gamma = (b_0, s_1, \dots, s_n, b_n, t, b_{n+1})$ with $\rho(\Gamma) = \gamma$ is equal to the number of chambers b_{n+1} such that $\delta(b_n, b_{n+1}) = t$ and $\rho(b_{n+1}) = c_{n+1}$.

Case 1: Suppose that $\delta(a, c_n)t > \delta(a, c_n)$. Thus $c_{n+1} \neq c_n$ (because γ is a -pointed) and hence $\delta(a, c_{n+1}) = \delta(a, c_n)t$. Since $c_n = \rho(b_n)$ we have $\delta(a, c_n) = \delta(a, b_n)$, and so $\delta(a, b_n)t > \delta(a, b_n)$. This implies that $\delta(a, b_{n+1}) = \delta(a, b_n)t$ for all b_{n+1} with $\delta(b_n, b_{n+1}) = t$, and therefore $\delta(a, \rho(b_{n+1})) = \delta(a, c_n)t = \delta(a, c_{n+1})$ and so $\rho(b_{n+1}) = c_{n+1}$. Thus there are q_t extensions Γ of Γ' so that $\rho(\Gamma) = \gamma$, and so

$$|\rho^{-1}(\gamma)| = |\rho^{-1}(\gamma')|q_t = q(\gamma')q_t.$$

We have $\alpha_t(\gamma) = \alpha_t(\gamma') + 1$, $\alpha_s(\gamma) = \alpha_s(\gamma')$ for all $s \neq t$, and $\sigma_s(\gamma) = \sigma_s(\gamma')$ for all $s \in S$, and so $q(\gamma')q_t = q(\gamma)$ completing the induction in this case.

Case 2: Suppose that $\delta(a, c_n)t < \delta(a, c_n)$ and that $c_n \neq c_{n+1}$, and so $\delta(a, c_{n+1}) = \delta(a, c_n)t$. Since $\rho(b_n) = c_n$ we have $\delta(a, b_n)t < \delta(a, b_n)$. Therefore $\rho(b_{n+1}) = c_{n+1}$ if and only if $\delta(a, b_{n+1}) = \delta(a, c_n)t = \delta(a, b_n)t$, and this occurs for $b_{n+1} \in \mathcal{C}_t(b_n)$ if and only if $b_{n+1} = \text{proj}_\pi(a)$ where π is the cotype t panel of b_n . Thus there is a unique extension Γ of Γ' so that $\rho(\Gamma) = \gamma$, and so

$$|\rho^{-1}(\gamma)| = |\rho^{-1}(\gamma')| = q(\gamma').$$

Since $\alpha_s(\gamma) = \alpha_s(\gamma')$ and $\sigma_s(\gamma) = \sigma_s(\gamma')$ for all $s \in S$ we have $q(\gamma') = q(\gamma)$.

Case 3: Suppose that $\delta(a, c_n)t < \delta(a, c_n)$ and that $c_n = c_{n+1}$. In this case $\rho(b_{n+1}) = c_{n+1}$ if and only if $\delta(a, b_{n+1}) = \delta(a, c_{n+1}) = \delta(a, c_n) = \delta(a, b_n)$, and this occurs for $b_{n+1} \in \mathcal{C}_t(b_n)$ if and only if $b_{n+1} \neq \text{proj}_\pi(b_n)$. Thus there are precisely $q_t - 1$ extensions Γ of Γ' so that $\rho(\Gamma) = \gamma$, and so

$$|\rho^{-1}(\gamma)| = |\rho^{-1}(\gamma')|(q_t - 1) = q(\gamma')(q_t - 1).$$

We have $\alpha_s(\gamma) = \alpha_s(\gamma')$ for all $s \in S$, $\sigma_s(\gamma) = \sigma_s(\gamma')$ for all $s \neq t$, and $\sigma_t(\gamma) = \sigma_t(\gamma') + 1$, and hence $q(\gamma) = q(\gamma')(q_t - 1)$ completing the proof. \square

A 1-pointed pregallery in the Coxeter complex $\Sigma(W, S)$ is simply called a *pointed pregallery*. Thus a pointed pregallery of type (s_1, \dots, s_n) is a sequence

$$\gamma = (w_0, s_1, w_1, s_2, \dots, s_n, w_n)$$

such that

1. $w_0, \dots, w_n \in W$ with $w_j \in \{w_{j-1}, w_{j-1}s_j\}$ for each $j = 1, \dots, n$, and
2. if $w_{j-1} = w_j$ then $\ell(w_{j-1}s_j) < \ell(w_{j-1})$.

If $w \in W$ and $\mathbf{v} = (s_1, \dots, s_n)$ let $\mathcal{P}(w, \mathbf{v})$ be the set of all pointed pregalleries in $\Sigma(W, S)$ of type \mathbf{v} starting at w , and for each $u \in W$ let

$$\mathcal{P}(w, \mathbf{v})_u = \{\gamma \in \mathcal{P}(w, \mathbf{v}) \mid \text{end}(\gamma) = u\}.$$

Theorem 3.6. *Let $u, v, w \in W$ and let $\mathbf{v} = (s_1, s_2, \dots, s_n)$ be a reduced expression for v . Then*

$$c_{u,v}^w = \sum_{\gamma \in \mathcal{P}(w, \mathbf{v})_u} q(\gamma).$$

Proof. Let $a, b \in \mathcal{C}(X)$ be chambers of X with $\delta(a, b) = w$. Let \mathcal{A} be an apartment of X containing a and b , and let $\rho = \rho_{\mathcal{A}, a}$. Since \mathbf{v} is reduced the map $\mathcal{G}(b, \mathbf{v}) \rightarrow \mathcal{C}_v(b)$, $\Gamma \mapsto \text{end}(\Gamma)$ is a bijection. Thus if c is the unique chamber of \mathcal{A} with $\delta(a, c) = u$ we have

$$|\mathcal{C}_u(a) \cap \mathcal{C}_v(b)| = |\{d \in \mathcal{C}_v(b) \mid \rho(d) = c\}| = |\{\Gamma \in \mathcal{G}(b, \mathbf{v}) \mid \rho(\text{end}(\Gamma)) = c\}|.$$

Thus by Proposition 3.5 we have

$$|\mathcal{C}_u(a) \cap \mathcal{C}_v(b)| = |\{\Gamma \in \rho^{-1}(c) \mid \Gamma \in \mathcal{P}_{\mathcal{A}, a}(b, \mathbf{v}), \text{end}(\Gamma) = c\}| = \sum_{\gamma \in \mathcal{P}_{\mathcal{A}, a}(b, \mathbf{v})_c} q(\gamma)$$

where $\mathcal{P}_{\mathcal{A}, a}(b, \mathbf{v})_c = \{\gamma \in \mathcal{P}_{\mathcal{A}, a}(b, \mathbf{v}) \mid \text{end}(\gamma) = c\}$. Since $\mathcal{A} \cong \Sigma(W, S)$ and $\delta(a, b) = w$ we may identify \mathcal{A} with $\Sigma(W, S)$ such that $a \leftrightarrow 1$ and $b \leftrightarrow w$. Under this identification, $\mathcal{P}_{\mathcal{A}, a}(b, \mathbf{v})_c$ becomes $\mathcal{P}(w, \mathbf{v})_u$, hence the result. \square

In the following theorem (the main theorem of this section) we present our combinatorial formula for the intersection numbers $c_{u,v}^w(I, J, K)$.

Theorem 3.7. *Let I, J, K be spherical subsets of S , and suppose that $u \in R(I, J)$, $v \in R(K, J)$, and $w \in R(I, K)$. Then*

$$c_{u,v}^w(I, J, K) = \sum_{\gamma} q(\gamma)$$

where the sum is over

$$\gamma \in \left\{ \begin{array}{l} \text{pointed pregalleries } \gamma \text{ of type } \mathbf{m}\mathbf{v} \text{ in } \Sigma(W, S) \\ \text{starting at } w \text{ and with } \text{end}(\gamma) \in W_I u W_J \end{array} \middle| m \in M_K(J, v) \right\},$$

where \mathbf{v} is a fixed reduced expression for v , and for each $m \in M_K(J, v)$ we fix a reduced expression \mathbf{m} .

Proof. This follows from Proposition 3.3 and Theorem 3.6. \square

Example 3.8. Let (W, S) be an F_4 Coxeter system with $S = \{1, 2, 3, 4\}$ in standard Bourbaki labelling and fix $I = \{2, 3, 4\}$. Consider an F_4 building with parameters $q_1 = q_2 = s$ and $q_3 = q_4 = t$. The minimal length double coset representatives for $W_I \backslash W / W_I$ are

$$w_0 = e, \quad w_1 = 1, \quad w_2 = 12321, \quad w_3 = 12324321, \quad w_4 = 123423121324321$$

where e is the identity element of W . In this example we apply Theorem 3.7 to compute the intersection cardinalities $c_{w_i, w_1}^{w_3}(I, I, I)$ for each $i = 0, 1, 2, 3, 4$. We have $W_I(I, w_1) = W_{\{3,4\}}$, and $M_I(I, w_1)$ consists of the 8 elements

$$m_0 = e, \quad m_1 = 2, \quad m_2 = 32, \quad m_3 = 432, \quad m_4 = 232, \quad m_5 = 2432, \quad m_6 = 32432, \quad m_7 = 232432.$$

To apply Theorem 3.7 we must find all pointed pregalleries of type $m_k w_1$ (with $0 \leq k \leq 7$) starting at w_3 , and compute $q(\gamma)$ and $\text{end}(\gamma)$ for each such pointed pregallery.

To abbreviate notation, we write a pointed pregallery γ of type $i_1 \cdots i_n$ starting at w as $\gamma = w \cdot \vec{i}_1 \vec{i}_2 \vec{i}_3 \widehat{i}_4 \cdots \vec{i}_n$ (for example), where the symbol \vec{i} indicates that the length increases in that step of the pregallery, \widehat{i} indicates length decreases in that step, and \widehat{i} indicates an i -stutter in the pregallery. Then $\text{end}(\gamma)$ is simply the word obtained by deleting all terms \widehat{i} .

Note that $\ell(w_j m_k) = \ell(w_j) + \ell(m_k)$ for all j, k , and so the first $\ell(m_k)$ steps of a pointed pregallery of type $m_k w_1$ starting at w_j are necessarily \vec{i} steps. Since $\ell(w_3 1) < \ell(w_3)$ we have two pointed pregalleries of type $m_0 w_1$ starting at w_3 , namely $\gamma_0 = w_3 \cdot \vec{1}$ and $\gamma'_0 = w_3 \cdot \widehat{1}$, and $q(\gamma_0) = 1$ and $q(\gamma'_0) = s - 1$. Similarly, for each $k = 1, 2, 3$ we have $\ell(w_3 m_k 1) < \ell(w_3 m_k)$, and so for each of these values of k there are two pointed pregalleries $\gamma_k = w_3 \cdot \vec{m}_k \vec{1}$ and $\gamma'_k = w_3 \cdot \vec{m}_k \widehat{1}$. We have $q(\gamma_1) = s$, $q(\gamma_2) = st$, $q(\gamma_3) = st^2$, $q(\gamma'_1) = (s - 1)s$, $q(\gamma'_2) = (s - 1)st$, and $q(\gamma'_3) = (s - 1)st^2$.

On the other hand, for $k = 4, 5, 6, 7$ we have $\ell(w_3 m_k 1) > \ell(w_3 m_k)$, and so for these values of k there is only 1 pointed pregallery of type $m_k w_1$ starting at w_3 , given by $\gamma_k = w_3 \cdot \vec{m}_k \vec{1}$. We have $q(\gamma_4) = s^3 t$, $q(\gamma_5) = s^3 t^2$, $q(\gamma_6) = s^3 t^3$, and $q(\gamma_7) = s^4 t^3$.

We see that $\text{end}(\gamma'_k) \in W_I w_3 W_I$ for all $k = 0, 1, 2, 3$, and $\text{end}(\gamma_0) \in W_I w_1 W_I$, and $\text{end}(\gamma_k) \in W_I w_2 W_I$ for $k = 1, 2, 3$, and $\text{end}(\gamma_k) \in W_I w_3 W_I$ for $k = 4, 5, 6$, and $\text{end}(\gamma_7) \in W_I w_4 W_I$. For example, consider γ_2 . Performing Coxeter moves gives $\text{end}(\gamma_2) = w_3 m_2 1 = 12324321321 = 4(12321)432 \in W_I w_2 W_I$.

Thus Theorem 3.7 gives $c_{w_0, w_1}^{w_3}(I, I, I) = 0$, $c_{w_1, w_1}^{w_3}(I, I, I) = 1$, and

$$\begin{aligned} c_{w_1, w_1}^{w_3}(I, I, I) &= 1, & c_{w_2, w_1}^{w_3}(I, I, I) &= s(t^2 + t + 1), \\ c_{w_3, w_1}^{w_3}(I, I, I) &= (s - 1)(st^2 + st + s + 1) + s^3 t(t^2 + t + 1), & c_{w_4, w_1}^{w_3}(I, I, I) &= s^4 t^3. \end{aligned}$$

Example 3.9. The formula in Theorem 3.7 is easily implemented in the MAGMA computational algebra system [5], making use of the existing Coxeter group package. For example, consider an E_8 Coxeter system with $S = \{1, \dots, 8\}$ in standard Bourbaki labelling and fix $I = S \setminus \{2\}$. Consider an E_8 building with parameter q . There are 35 distinct $W_I \backslash W / W_I$ double cosets. Let w_i , with $0 \leq i \leq 34$, denote the minimal length representatives of these double cosets. Fixing an order, we take $w_0 = e$, $w_1 = 2$, $w_2 = 243542$, $w_3 = 24315436542$, $w_4 = 243542654376542$, $w_5 = 2431542654376542$, and $w_6 = 2435426543176543876542$. Write $C_{i,j}^k = c_{w_i, w_j}^{w_k}(I, I, I)$. Implementing Theorem 3.7 into MAGMA we obtain

$$\begin{aligned} C_{2,1}^1 &= \phi_2(q^2) \phi_3(q) \phi_5(q) q^5 & C_{2,1}^2 &= \phi_2(q^2) (\phi_3(q)^2 q^3 - 1) & C_{2,1}^3 &= \phi_3(q)^2 q \\ C_{2,1}^4 &= \phi_3(q^2) \phi_5(q) & C_{2,1}^5 &= \phi_3(q) & C_{2,1}^6 &= 1, \end{aligned}$$

where $\phi_n(x) = x^{n-1} + \cdots + x + 1$ and $C_{2,1}^k = 0$ for $k = 0$ and $7 \leq k \leq 34$.

4 Hecke operators on regular buildings

If Γ is a distance regular graph with intersection numbers $a_{k,\ell}^m$ then, as discussed in the introduction, the k -adjacency operators

$$A_k f(x) = \sum_{y \in F_k(x)} f(y) \quad \text{for } f : V \rightarrow \mathbb{C}$$

satisfy linear relations

$$A_k A_\ell = \sum_{m \in \mathbb{N}} a_{k,\ell}^m A_m \quad \text{for all } k, \ell \in \mathbb{N}.$$

Thus the vector space over \mathbb{C} with basis $\{A_k \mid k \in \mathbb{N}\}$ is an algebra \mathcal{A} .

This brings us to an investigation of generalised adjacency operators on buildings, one of our primary motivations for studying intersection cardinalities of generalised spheres in regular buildings. For each pair I, J of spherical subsets, and each $w \in R(I, J)$, define the (I, J, w) -adjacency operator $T_w^{IJ} : \{f : X_J \rightarrow \mathbb{C}\} \rightarrow \{f : X_I \rightarrow \mathbb{C}\}$ by

$$(T_w^{IJ} f)(A) = \sum_{B \in F(A, J, w)} f(B), \quad \text{for all } A \in X_I \text{ and } f : X_J \rightarrow \mathbb{C}.$$

We also refer to these operators as *Hecke operators* on the building.

Proposition 4.1. *Let I, J and K be spherical, and suppose that $u \in R(I, J)$ and $v \in R(K, J)$. Then*

$$T_u^{IJ} T_v^{JK} = \sum_{w \in R(I, K)} c_{u, v^{-1}}^w(I, J, K) T_w^{IK},$$

where for fixed u and v , $c_{u, v^{-1}}^w(I, J, K)$ is nonzero for only finitely many $w \in R(I, K)$.

Proof. Directly from definition of the adjacency operators we compute, for $A \in X_I$,

$$\begin{aligned} (T_u^{IJ} T_v^{JK} f)(A) &= \sum_{B \in F(A, J, u)} \sum_{D \in F(B, K, v)} f(D) \\ &= \sum_{D \in X_K} \sum_{B \in X_J} 1_{F(A, J, u)}(B) 1_{F(B, K, v)}(D) f(D), \end{aligned}$$

where for any subset $F \subseteq X$ we write 1_F for the indicator function of F . Since $D \in F(B, K, v)$ if and only if $B \in F(D, J, v^{-1})$ we have $1_{F(B, K, v)}(D) = 1_{F(D, J, v^{-1})}(B)$, and thus

$$(T_u^{IJ} T_v^{JK} f)(A) = \sum_{D \in X_K} |F(A, J, u) \cap F(D, J, v^{-1})| f(D).$$

Since X_K is the disjoint union of the sets $F(A, K, w)$ with $w \in R(I, K)$ we have

$$(T_u^{IJ} T_v^{JK} f)(A) = \sum_{w \in R(I, K)} \sum_{D \in F(A, K, w)} |F(A, J, u) \cap F(D, J, v^{-1})| f(D),$$

and the result follows from Theorem 3.1 and Remark 3.2. \square

Corollary 4.2. *Let I and J be spherical, and let $\mathcal{A}(I, J)$ be the vector space over \mathbb{C} spanned by the operators $\{T_w^{IJ} \mid w \in R(I, J)\}$. Then $\{T_w^{IJ} \mid w \in R(I, J)\}$ is a basis of $\mathcal{A}(I, J)$. Moreover, $\mathcal{A}(I, I)$ is an algebra, and $\mathcal{A}(I, J)$ is a left $\mathcal{A}(I, I)$ -module and a right $\mathcal{A}(J, J)$ -module.*

Proof. Let $T = \sum_{w \in R(I, J)} a_w T_w^{IJ}$, with $a_w \in \mathbb{C}$. Let $A \in X_I$, and $B \in F(A, J, v)$ with $v \in R(I, J)$, and let $\delta_B : X_J \rightarrow \mathbb{C}$ be the function $\delta_B(B) = 1$ and $\delta_B(B') = 0$ for all $B' \in X_J \setminus \{B\}$. Then $T\delta_B(A) = a_v$, and hence the operators $\{T_w^{IJ} \mid w \in R(I, J)\}$ are linearly independent, and thus form a basis of $\mathcal{A}(I, J)$. Proposition 4.1 shows that $\mathcal{A}(I, I)$ is closed under multiplication, and that $\mathcal{A}(I, J)$ is closed under left multiplication by elements of $\mathcal{A}(I, I)$ and right multiplication by elements of $\mathcal{A}(J, J)$. Hence the result. \square

In the language of *association schemes* (see [8]), the algebra $\mathcal{A}(I, I)$ is the *Bose-Mesner algebra* of the natural association scheme on the cotype I simplices of X .

Example 4.3. Consider the F_4 example of Example 3.8. Write $A_i = T_{w_i}^{II}$ for $i = 0, 1, 2, 3, 4$. Note that the minimal length double coset representatives w_i are involutions. This implies that $\mathcal{A}(I, I)$ is commutative, for if $\delta(A, B) = w \in R(I, I)$ then $\delta(B, A) = w^{-1} = w$ and so

$$c_{u,v^{-1}}^w(I, I, I) = |F(A, I, u) \cap F(B, I, v^{-1})| = |F(B, I, v) \cap F(A, I, u^{-1})| = c_{v,u^{-1}}^w(I, I, I).$$

By Proposition 4.1, the computations in Example 3.8 give the coefficient of A_3 in the expansions of A_1A_1 , A_2A_1 , A_3A_1 , and A_4A_1 . We obtain:

$$\begin{aligned} A_1A_1 &= s(s+1)(st+1)(st^2+1)A_0 + [s^2(t^2+t+1) + s-1]A_1 + (s+1)(st+1)A_2 + A_3 \\ A_2A_1 &= s^3t(t^2+t+1)A_1 + (s^2-1)(st+1)A_2 + s(t^2+t+1)A_3 \\ A_3A_1 &= s^4t^3A_1 + s^2t^2(s+1)(st+1)A_2 + [(s-1)(st^2+st+s+1) + s^3t(t^2+t+1)]A_3 \\ &\quad + (s+1)(st+1)(st^2+1)A_4 \\ A_4A_1 &= s^4t^3A_3 + (s^2-1)(st+1)(st^2+1)A_4. \end{aligned}$$

We note that these formulae are sufficient to compute the character table for the algebra $\mathcal{A}(I, I)$.

Example 4.4. Consider the E_8 example of Example 3.9. Write $A_i = T_{w_i}^{II}$ for $0 \leq i \leq 34$. In this case the algebra spanned by the operators A_i is not commutative, for example in the notation of Example 3.9 we have $C_{2,1}^6 = 1$ and $C_{1,2}^6 = 0$ (see Corollary 5.2 for more on commutativity). Proposition 4.1 gives $A_2A_1 = C_{2,1}^1A_1 + C_{2,1}^2A_2 + C_{2,1}^3A_3 + C_{2,1}^4A_4 + C_{2,1}^5A_5 + C_{2,1}^6A_6$.

Consider the case $I = J = \emptyset$. Write $T_w^{\emptyset\emptyset} = T_w$ for all $w \in W$ and let $\mathcal{H} = \mathcal{A}(\emptyset, \emptyset)$. By Proposition 4.1 we have

$$T_uT_v = \sum_{w \in W} c_{u,v^{-1}}^w T_w \quad \text{for all } u, v \in W, \quad (4.1)$$

where the numbers $c_{u,v^{-1}}^w$ are defined in (3.1). From the geometry of the building it is not hard to show that (see [21, Theorem 3.4])

$$T_wT_s = \begin{cases} T_{ws} & \text{if } \ell(ws) > \ell(w) \\ q_s T_{ws} + (q_s - 1)T_w & \text{if } \ell(ws) < \ell(w). \end{cases} \quad (4.2)$$

Thus the algebra \mathcal{H} is isomorphic to the extensively studied (*Iwahori*) *Hecke algebra* of the Coxeter system (W, S) (see [16]). Our goal now is to interpret our more general algebras and modules $\mathcal{A}(I, J)$ in terms of the more familiar Hecke algebra \mathcal{H} .

For each spherical subset I of S define an element $\mathbf{1}_I$ of \mathcal{H} by

$$\mathbf{1}_I = \frac{1}{N(I)} \sum_{w \in W_I} T_w.$$

The subalgebra $\mathbf{1}_I \mathcal{H} \mathbf{1}_I$ of \mathcal{H} is called a *parabolic Hecke algebra* (see [3, 11]). Note that $\mathbf{1}_I \mathcal{H} \mathbf{1}_I$ and \mathcal{H} have different units ($\mathbf{1}_I$ and T_1 respectively).

Lemma 4.5. *Let I be spherical. Then $T_u \mathbf{1}_I = \mathbf{1}_I T_u = q_u \mathbf{1}_I$ for all $u \in W_I$, and $\mathbf{1}_I^2 = \mathbf{1}_I$.*

Proof. By (4.2) the vector space spanned by $\{T_u \mid u \in W_I\}$ is a subalgebra of \mathcal{H} . Therefore if $u \in W_I$ we have

$$T_u \mathbf{1}_I = \frac{1}{N(I)} \sum_{v \in W_I} T_u T_v = \frac{1}{N(I)} \sum_{w \in W_I} \left(\sum_{v \in W_I} c_{u,v^{-1}}^w \right) T_w.$$

If $w \in W_I$ and $a, b \in \mathcal{C}$ with $b \in \mathcal{C}_w(a)$ then $\sum_v c_{u,v}^w = \sum_v |\mathcal{C}_u(a) \cap \mathcal{C}_{v^{-1}}(b)| = |\mathcal{C}_u(a)| = q_u$. Thus $T_u \mathbf{1}_I = q_u \mathbf{1}_I$, and the proof that $\mathbf{1}_I T_u = q_u \mathbf{1}_I$ is similar.

Then

$$\mathbf{1}_I^2 = \frac{1}{N(I)} \sum_{w \in W_I} T_w \mathbf{1}_I = \frac{1}{N(I)} \sum_{w \in W_I} q_w \mathbf{1}_I = \frac{N(I)}{N(I)} \mathbf{1}_I = \mathbf{1}_I,$$

completing the proof. \square

For spherical subsets I and J of S and $w \in R(I, J)$ define an element $P_w^{IJ} \in \mathbf{1}_I \mathcal{H} \mathbf{1}_J$ by

$$P_w^{IJ} = \frac{N(I)}{N(I \cap wJw^{-1})} \mathbf{1}_I T_w \mathbf{1}_J.$$

The reason for this choice of normalisation will become clear in Theorem 4.9.

Lemma 4.6. *If I and J are spherical subsets of S and $w \in R(I, J)$ then*

$$P_w^{IJ} = \frac{1}{N(J)} \sum_{z \in W_I w W_J} T_z.$$

Proof. Using (1.2)

$$\mathbf{1}_I T_w \mathbf{1}_J = \frac{1}{N(I)} \sum_{u \in W_I} T_u T_w \mathbf{1}_J = \frac{1}{N(I)} \sum_{x \in M_I(J, w)} \sum_{y \in W_I(J, w)} T_x T_y T_w \mathbf{1}_J.$$

Since w is (I, J) -reduced, and since $y \in W_I(J, w)$, we have $\ell(yw) = \ell(y) + \ell(w)$. Moreover $yw = wy'$ with $y' = w^{-1}yw \in W_J$ and $\ell(wy') = \ell(w) + \ell(y')$. This implies that $q_{y'} = q_y$ (because $q_y q_w = q_{yw} = q_{wy'} = q_w q_{y'}$), and by (4.2) and Lemma 4.5 we have

$$T_y T_w \mathbf{1}_J = T_{yw} \mathbf{1}_J = T_{wy'} \mathbf{1}_J = T_w T_{y'} \mathbf{1}_J = q_{y'} T_w \mathbf{1}_J = q_y T_w \mathbf{1}_J.$$

Thus by (1.1) we have $\sum_{y \in W_I(J, w)} T_x T_y T_w \mathbf{1}_J = N(I \cap wJw^{-1}) T_x T_w \mathbf{1}_J$, and so using Lemma 1.1,

$$P_w^{IJ} = \frac{N(I)}{N(I \cap wJw^{-1})} \mathbf{1}_I T_w \mathbf{1}_J = \sum_{x \in M_I(J, w)} T_x T_w \mathbf{1}_J = \frac{1}{N(J)} \sum_{z \in W_I w W_J} T_z. \quad \square$$

Remark 4.7. Lemma 4.6 gives an independent algebraic proof of Theorem 2.1, as follows. The linear map $\pi : \mathcal{H} \rightarrow \mathbb{C}$ with $\pi(T_w) = q_w$ for all $w \in W$ satisfies

$$\pi(T_w) \pi(T_s) = \begin{cases} \pi(T_{ws}) & \text{if } \ell(ws) > \ell(w) \\ q_s \pi(T_{ws}) + (q_s - 1) \pi(T_w) & \text{if } \ell(ws) < \ell(w), \end{cases}$$

and so by (4.2) the map π is a 1-dimensional representation of \mathcal{H} . Applying this representation to the formula in Lemma 4.6 gives, for $w \in R(I, J)$,

$$\frac{N(I)}{N(I \cap wJw^{-1})} q_w = \frac{1}{N(J)} \sum_{z \in W_I w W_J} q_z. \quad (4.3)$$

If $A \in X_I$ then $\mathcal{C}(F(A, J, w)) = \bigsqcup_{z \in W_I w W_J} \mathcal{C}_z(a)$ for any fixed $a \in \mathcal{C}(A)$. Thus

$$|\mathcal{C}(F(A, J, w))| = \sum_{z \in W_I w W_J} q_z,$$

and since each simplex $B \in F(A, J, w)$ is contained in $N(J)$ chambers of $\mathcal{C}(F(A, J, w))$, the right hand side of (4.3) equals $|F(A, J, w)|$, proving Theorem 2.1.

Corollary 4.8. *If I and J are spherical subsets of S then the vector space $\mathbf{1}_I \mathcal{H} \mathbf{1}_J$ has basis $\{P_w^{IJ} \mid w \in R(I, J)\}$.*

Proof. Let $z \in W$. Then $z \in W_I w W_J$ for some $w \in R(I, J)$, and we can write $z = xwy$ with $x \in M_I(J, w)$ and $y \in W_J$. Thus $\ell(z) = \ell(x) + \ell(w) + \ell(y)$. Using (4.2) and Lemma 4.5 we have

$$\mathbf{1}_I T_z \mathbf{1}_J = \mathbf{1}_I T_x T_w T_y \mathbf{1}_J = q_x q_y \mathbf{1}_I T_w \mathbf{1}_J.$$

Hence the set $\{P_w^{IJ} \mid w \in R(I, J)\}$ spans $\mathbf{1}_I \mathcal{H} \mathbf{1}_J$. It follows from Lemma 4.6 and the linear independence of the T_z , $z \in W$, that the operators P_w^{IJ} , $w \in R(I, J)$, are linearly independent. \square

The following theorem, which shows that the multiplication table for the operators T_w^{IJ} and P_w^{IJ} are the same, is the main result of this section.

Theorem 4.9. *Let I, J, K be spherical, and suppose that $u \in R(I, J)$ and $v \in R(K, J)$. Then*

$$P_u^{IJ} P_v^{JK} = \sum_{w \in R(I, K)} c_{u, v^{-1}}^w(I, J, K) P_w^{IK}$$

where $c_{u, v^{-1}}^w(I, J, K)$ is as in Theorem 3.1. In particular, the linear map

$$\theta : \begin{array}{ccc} \mathcal{A}(I, I) & \longrightarrow & \mathbf{1}_I \mathcal{H} \mathbf{1}_I \\ T_w^{II} & \longmapsto & P_w^{II} \end{array} \quad \text{is an isomorphism of algebras,}$$

and $\mathcal{A}(I, J)$ is a left $\mathbf{1}_I \mathcal{H} \mathbf{1}_I$ -module and a right $\mathbf{1}_J \mathcal{H} \mathbf{1}_J$ -module.

Proof. By Lemma 4.6, the expansion $T_x T_y = \sum_{z \in W} c_{x, y^{-1}}^z T_z$ (see (4.1)), and the decomposition of W into $W_I w W_K$ double cosets with $w \in R(I, K)$, we have

$$\begin{aligned} P_u^{IJ} P_v^{JK} &= \frac{1}{N(J)N(K)} \sum_{\substack{x \in W_I u W_J \\ y \in W_J v W_K}} T_x T_y \\ &= \frac{1}{N(J)N(K)} \sum_{w \in R(I, K)} \sum_{z \in W_I w W_K} \left(\sum_{\substack{x \in W_I u W_J \\ y \in W_J v W_K}} c_{x, y^{-1}}^z \right) T_z. \end{aligned}$$

Thus by Theorem 3.1 and Lemma 4.6 we have

$$P_u^{IJ} P_v^{JK} = \sum_{w \in R(I, K)} \frac{c_{u, v^{-1}}^w(I, J, K)}{N(K)} \sum_{z \in W_I w W_K} T_z = \sum_{w \in R(I, K)} c_{u, v^{-1}}^w(I, J, K) P_w^{IK}.$$

It follows that the linear map $\theta : \mathcal{A}(I, I) \rightarrow \mathbf{1}_I \mathcal{H} \mathbf{1}_I$ with $T_w^{II} \mapsto P_w^{II}$ is an isomorphism of algebras by using the fact that $\mathcal{A}(I, I)$ has basis $\{T_w^{II} \mid w \in R(I, I)\}$ and $\mathbf{1}_I \mathcal{H} \mathbf{1}_I$ has basis $\{P_w^{II} \mid w \in R(I, I)\}$. The final statement follows from Corollary 4.2. \square

5 Applications

In this final section we describe some applications of our main theorems.

5.1 Structure constants in parabolic Hecke algebras

Theorem 4.9 gives a combinatorial interpretation of the structure constants in parabolic Hecke algebras, and Theorem 3.7 gives a formula for these structure constants. In the case when W has type \tilde{A}_n and $I = S \setminus \{s_0\}$ these structure constants are *Hall polynomials*, and nonnegativity of these polynomials was proved by Miller Malley [20]. More generally, if W is of affine type and $I = S \setminus \{s_0\}$ then the structure constants are the structure constants of the algebra spanned by the *Macdonald spherical functions*, and nonnegativity of these structure constants was proved independently by Parkinson [21] and Schwer [28]. Our formula in Theorem 3.7 immediately gives a considerable generalisation of these results to arbitrary type:

Corollary 5.1. *The structure constants in the parabolic Hecke algebra $\mathbf{1}_I \mathcal{H} \mathbf{1}_I$ relative to the basis $\{P_w^{II} \mid w \in R(I, I)\}$ are nonnegative.*

5.2 Commutativity of algebras of Hecke operators on buildings

Recall that for distance regular graphs the algebra \mathcal{A} of averaging operators is necessarily commutative (this boils down to the fact that the graph distance satisfies $d(x, y) = d(y, x)$). In contrast, the algebra $\mathcal{A}(I, I)$ is not necessarily commutative. Using Theorem 4.9 we have the following commutativity classification (where we use standard Bourbaki labelling [6]), showing that in fact commutativity is extremely rare.

Corollary 5.2. *Suppose that (W, S) is irreducible. Let I be a spherical subset of S . The algebra $\mathcal{A}(I, I)$ is noncommutative if $|S \setminus I| > 1$. If $I = S \setminus \{i\}$ then $\mathcal{A}(I, I)$ is commutative in the cases*

1. $W = A_n$ and $1 \leq i \leq n$, $W = B_n$ and $1 \leq i \leq n$, $W = D_n$ and $1 \leq i \leq n/2$ or $i = n - 1, n$, $W = E_6$ and $i = 1, 2, 6$, $W = E_7$ and $i = 1, 2, 7$, $W = E_8$ and $i = 1, 8$, $W = F_4$ and $i = 1, 4$, $W = H_3$ and $i = 1, 3$, $W = H_4$ and $i = 1$, $W = I_2(p)$ and $i = 1, 2$, or
2. W affine and i is a special type (that is, i is in the orbit of the special node 0 under the action of diagram automorphisms),

and noncommutative otherwise.

Proof. By Theorem 4.9 the algebra $\mathcal{A}(I, I)$ is isomorphic to the parabolic Hecke algebra $\mathbf{1}_I \mathcal{H} \mathbf{1}_I$. The commutative parabolic Hecke algebras are classified in [3, Theorem 2.1], and the result follows from this classification. \square

5.3 Combinatorics of double cosets in groups with BN -pairs

Let G be a group acting on X by type preserving simplicial automorphisms. The group G has a *strongly transitive* action on X (with respect to some fixed system of apartments) if G is transitive on the set of pairs (\mathcal{A}, a) with \mathcal{A} an apartment and $a \in \mathcal{A}$ a chamber of \mathcal{A} . The group G has a *Weyl-transitive* action on X if for all $w \in W$ the action is transitive on all ordered pairs (a, b) of chambers such that $\delta(a, b) = w$. Recall that strong transitivity implies Weyl transitivity, although the reverse implication does not hold in general (see [1]).

If G acts Weyl transitively on X then it is immediate that $|F(A, J, u) \cap F(A', J, v)|$ depends only on u, v and $\delta(A, A')$, and therefore Theorem 3.1 is most interesting in that case where the building does not admit a Weyl transitive group action.

We now outline applications of our formula in Theorem 3.7 to double coset combinatorics in groups G acting strongly transitively on X . Fix a chamber $a \in \mathcal{C}$ and an apartment \mathcal{A} with

$a \in \mathcal{A}$. Let $B = \text{stab}_G(a)$ and $N = \text{stab}_G(\mathcal{A})$. The pointwise stabiliser of \mathcal{A} is $H = B \cap N$, and N/H is isomorphic to W (the Coxeter group of X). It is well-known that (G, B, N, W) is a BN -pair (also called a Tits system, see [2, Theorem 6.56]). The stabiliser of the cotype I simplex $A = a_I$ of a is the *parabolic subgroup* $P_I = \bigsqcup_{w \in W_I} BwB$.

Proposition 5.3. *With the above notation, if I, J and K are spherical, $u \in R(I, J)$, $v \in R(K, J)$, and $w \in R(I, K)$, then*

$$\text{Card}_{G/P_J}(P_I u P_J \cap w P_K v P_J) = c_{u,v}^w(I, J, K).$$

Thus Theorem 3.7 gives a formula for these cardinalities.

Proof. The cotype I simplices X_I of X can be identified with G/P_I . Let $A = P_I$ under this identification. If $w \in R(I, J)$ then $F(A, J, w)$ is the set of P_J cosets in $P_I w P_J$. Therefore if I, J and K are spherical, and if $u \in R(I, J)$, $v \in R(K, J)$, and $w \in R(I, K)$, then

$$\text{Card}_{G/P_J}(P_I u P_J \cap w P_K v P_J) = |F(A, J, u) \cap F(wA, J, v)| = c_{u,v}^w(I, J, K). \quad \square$$

5.4 Convolution algebras and Gelfand pairs

Suppose that G is a locally compact group acting strongly transitively on the building X , and let $B = \text{stab}_G(a)$ be the stabiliser of a fixed chamber (as in the previous section). Assume that B is compact. A function $f : G \rightarrow \mathbb{C}$ is *bi- B -invariant* if $f(bgb') = f(g)$ for all $g \in G$ and all $b, b' \in B$. Let $\mathcal{F}(B \backslash G/B)$ be the space of bi- B -invariant complex valued functions supported on finitely many double cosets. Define convolution in $\mathcal{F}(B \backslash G/B)$ by

$$(f_1 * f_2)(g) = \int_G f_1(gh) f_2(h^{-1}) d\mu(h),$$

where μ is a Haar measure on G normalised so that $\mu(B) = 1$. It is elementary that $\mathcal{F}(B \backslash G/B)$ is closed under convolution, and so $\mathcal{F}(B \backslash G/B)$ is an algebra.

For $I, J \subseteq S$ spherical and $w \in R(I, J)$, let $\psi_{P_I w P_J} = \frac{1}{N(J)} \chi_{P_I w P_J}$, where $\chi_{P_I w P_J}$ is the characteristic function of the double coset $P_I w P_J$.

Proposition 5.4. *Let $I, J, K \subseteq S$ be spherical, and let $u \in R(I, J)$ and $v \in R(J, K)$. Then*

$$\psi_{P_I u P_J} * \psi_{P_J v P_K} = \sum_{w \in R(I, K)} c_{u,v^{-1}}^w(I, J, K) \psi_{P_I w P_K}.$$

Proof. The assumption that I and J are spherical ensures that $\psi_{P_I w P_J}$ is supported on finitely many B double cosets, thus $\psi_{P_I w P_J} \in \mathcal{F}(B \backslash G/B)$. If $g_1 \in P_I$ and $g_2 \in P_K$ then for all $g \in G$ we have

$$\begin{aligned} (\psi_{P_I u P_J} * \psi_{P_J v P_K})(g_1 g g_2) &= \int_G \psi_{P_I u P_J}(g_1 g g_2 h) \psi_{P_J v P_K}(h^{-1}) d\mu(h) \\ &= \int_G \psi_{P_I u P_J}(gh') \psi_{P_J v P_K}(h'^{-1} g_2) d\mu(h') = (\psi_{P_I u P_J} * \psi_{P_J v P_K})(g). \end{aligned}$$

Thus $\psi_{P_I u P_J} * \psi_{P_J v P_K}$ is left P_I -invariant and right P_K -invariant, and hence is a linear combination of terms $\psi_{P_I w P_K}$ with $w \in R(I, K)$. To compute the coefficient of $\psi_{P_I w P_K}$ in this linear

combination, note that from the definition of convolution, and using Proposition 5.3, we have

$$\begin{aligned} \frac{1}{N(J)}(\psi_{P_I u P_J} * \psi_{P_J v P_J})(w) &= \frac{1}{N(J)} \text{Card}_{G/B}(w^{-1} P_I u P_J \cap P_K v^{-1} P_J) \\ &= \frac{1}{N(J)} \text{Card}_{G/B}(P_I u P_J \cap w P_K v^{-1} P_J) \\ &= \text{Card}_{G/P_J}(P_I u P_J \cap w P_K v^{-1} P_J) = c_{u,v^{-1}}^w(I, J, K), \end{aligned}$$

completing the proof. \square

Corollary 5.5. *With the above notation, let $\mathcal{F}(P_I \backslash G / P_I)$ be the subalgebra of $\mathcal{F}(B \backslash G / B)$ consisting of bi- P_I -invariant functions. Then*

$$\mathcal{F}(P_I \backslash G / P_I) \cong \mathbf{1}_I \mathcal{H} \mathbf{1}_I,$$

with the isomorphism given by $\psi_{P_I w P_I} \mapsto P_w^{II}$ for $w \in R(I, I)$.

Proof. This is immediate from Theorem 4.9 and Proposition 5.4. \square

Recall that for any locally compact group G and compact subgroup K , the pair (G, K) is called a *Gelfand pair* if the convolution algebra of compactly supported continuous bi- K -invariant functions on G is commutative.

Corollary 5.6. *The pair (G, P_I) from Corollary 5.5 is a Gelfand pair if and only if $\mathcal{A}(I, I)$ is commutative, and hence only in the cases listed in Corollary 5.2.*

Proof. From Theorem 4.9 and Corollary 5.5 we have $\mathcal{A}(I, I) \cong \mathbf{1}_I \mathcal{H} \mathbf{1}_I \cong \mathcal{F}(P_I \backslash G / P_I)$ and the result follows. \square

Note that Corollary 5.6 implies that the only Gelfand pairs arising from buildings with infinite Coxeter groups are those Gelfand pairs (G, K) where K is the stabiliser of a special vertex in an affine building. These Gelfand pairs have received considerable attention (see for example [9]).

5.5 Isotropic random walks on the simplices of buildings

Fix $I \subseteq S$ spherical, and let $(Z_n)_{n \geq 0}$ be a random walk on the set X_I of all cotype I simplices of a locally finite regular building X . Thus $(Z_n)_{n \geq 0}$ is a Markov chain of X_I -valued random variables, with evolution governed by *transition probabilities*

$$p(A, B) = \mathbb{P}[Z_{n+1} = B \mid Z_n = A] \quad \text{for } A, B \in X_I.$$

For simplicity we assume throughout that $(Z_n)_{n \geq 0}$ has *bounded jumps*. That is, for each $A \in X_I$ the probability $p(A, B)$ is nonzero for only finitely many $B \in X_I$.

One important goal in random walk theory is to obtain estimates for the n -step transition probabilities

$$p^{(n)}(A, B) = \mathbb{P}[Z_n = B \mid Z_0 = A],$$

and in particular for the n -step return probabilities $p^{(n)}(A, A)$. Such a result is called a *local limit theorem*, and ideally it takes the form of an asymptotic formula.

A random walk on X_I is called *isotropic* if $p(A, B) = p(A', B')$ whenever $\delta(A, B) = \delta(A', B')$. In other words, a walk is isotropic if the transition probabilities depend only on the Weyl distance.

In this section we use Theorem 4.9 to reduce the ‘general’ case of isotropic random walks on the cotype I simplices of a building to the ‘special’ case of isotropic random walks on the set of chambers of the building (where $I = \emptyset$). This latter case admits a rather complete theory when X is an affine building (see [25]), and thus we can now obtain precise local limit theorems for isotropic random walks on the cotype I simplices of an affine building. We will give a concrete example at the end of this section.

The *transition operator* of a random walk $(Z_n)_{n \geq 0}$ on X_I is the operator T acting on the space of all functions $f : X_I \rightarrow \mathbb{C}$ by

$$Tf(A) = \sum_{B \in X_I} p(A, B)f(B).$$

Thus the transition operator of an isotropic random walk $(Z_n)_{n \geq 0}$ on X_I is given by

$$T = \sum_{w \in R(I, I)} p_w T_w^{II} \quad \text{where } p_w = p(A, B) \text{ for any } A, B \in X_I \text{ with } \delta(A, B) = w, \quad (5.1)$$

and so T is an element of the algebra $\mathcal{A}(I, I)$ (here we use the bounded jumps assumption to ensure that T is a finite linear combination; in the general case T lies in a completion of the algebra $\mathcal{A}(I, I)$). It is useful to note that if $\delta(A, B) = w$ then $p^{(n)}(A, B)$ is the coefficient of T_w^{II} in the expansion of $T^n \in \mathcal{A}(I, I)$ as a linear combination of the basis $\{T_v^{II} \mid v \in R(I, I)\}$.

Proposition 5.7. *Let $I \subseteq S$ be spherical. Let $(Z_n)_{n \geq 0}$ be an isotropic random walk on X_I with bounded jumps and transition probabilities $p(A, B)$, and let $p_w = p(A, B)$ for any $A, B \in X_I$ with $\delta(A, B) = w$. Let $(\bar{Z}_n)_{n \geq 0}$ be the isotropic random walk on \mathcal{C} with transition probabilities*

$$\bar{p}(a, b) = \frac{p_w}{N(I)} \quad \text{if } \delta(a, b) \in W_I w W_I \text{ with } w \in R(I, I).$$

Then for all $A, B \in X_I$ and all $a, b \in \mathcal{C}$ with $\delta(a, b) \in W_I \delta(A, B) W_I$ we have

$$p^{(n)}(A, B) = \frac{1}{N(I)} \bar{p}^{(n)}(a, b) \quad \text{for all } n \in \mathbb{N}.$$

Proof. We must first check that $\bar{p}(a, b)$ defines a transition probability, and thus we verify that $\sum_{b \in \mathcal{C}} \bar{p}(a, b) = 1$ for each $a \in \mathcal{C}$. We have

$$\sum_{b \in \mathcal{C}} \bar{p}(a, b) = \sum_{w \in R(I, I)} \sum_{\{b \mid \delta(a, b) \in W_I w W_I\}} \bar{p}(a, b) = \sum_{w \in R(I, I)} \frac{p_w}{N(I)} |\{b \mid \delta(a, b) \in W_I w W_I\}|.$$

The cardinality in the summand is equal to $N(I)|F(A, I, w)|$ where A is the cotype I simplex of a , since each element of $F(A, I, w)$ is contained in exactly $N(I)$ chambers of $\{b \mid \delta(a, b) \in W_I w W_I\}$. Thus

$$\sum_{b \in \mathcal{C}} \bar{p}(a, b) = \sum_{w \in R(I, I)} p_w |F(A, I, w)| = \sum_{B \in X_I} p(A, B) = 1.$$

Let $T \in \mathcal{A}(I, I)$ be the transition operator of $(Z_n)_{n \geq 0}$, as in (5.1). Let $\theta : \mathcal{A}(I, I) \rightarrow \mathbf{1}_I \mathcal{H} \mathbf{1}_I$ be the isomorphism from Theorem 4.9. By Lemma 4.6

$$\theta(T) = \sum_{w \in R(I, I)} p_w P_w^{II} = \sum_{w \in R(I, I)} \frac{p_w}{N(I)} \sum_{z \in W_I w W_I} T_z = \sum_{z \in W} \bar{p}_z T_z,$$

where $\bar{p}_z = p_w/N(I)$ if $z \in W_I w W_I$ with $w \in R(I, I)$. Thus $\theta(T) \in \mathcal{H}$ is the transition operator of the isotropic random walk $(\bar{Z})_{n \geq 0}$ on \mathcal{C} .

Let $A, B \in X_I$ with $\delta(A, B) = w \in R(I, I)$. Then $p^{(n)}(A, B)$ is the coefficient of T_w^{II} in the expansion of T^n , and this equals the coefficient of P_w^{II} in the expansion of $\theta(T^n) = \theta(T)^n$. By Lemma 4.6 this equals $N(I)^{-1}$ times the coefficient of T_z in the expansion of $\theta(T)^n$ for any $z \in W_I w W_I$, and this equals $\bar{p}^{(n)}(a, b)/N(I)$ for any $a, b \in \mathcal{C}$ with $\delta(a, b) = z \in W_I w W_I$. \square

We conclude with a concrete example illustrating how Proposition 5.7 can be used, in conjunction with the techniques in [25], to give new local limit theorems for random walks on simplices of affine buildings.

Let X be a locally finite thick \tilde{A}_2 building with thickness parameter q . Let the vertices of X have types $\{0, 1, 2\}$ and let X_0 be the set of all cotype 0 simplices of X (that is, the cotype 0 panels of X). The ‘neighbours’ of $A \in X_0$ are those simplices $B \in X_0$ with $d(A, B) = 1$, or equivalently, $\delta(A, B) \in \{s_1, s_2\}$. Note that $|\{B \in X_0 \mid d(A, B) = 1\}| = 2q(q+1)$ for each $A \in X_0$. The *simple random walk* on X_0 is the random walk with transition probabilities

$$p(A, B) = \begin{cases} \frac{1}{2q(q+1)} & \text{if } d(A, B) = 1 \\ 0 & \text{otherwise.} \end{cases}$$

This walk is isotropic, with transition operator

$$T = \frac{1}{2q(q+1)} (T_{s_1}^{00} + T_{s_2}^{00}).$$

Let $\bar{p}(a, b)$ be the transition probabilities of the associated isotropic random walk $(\bar{Z}_n)_{n \geq 0}$ on \mathcal{C} (as in Proposition 5.7). The transition operator of this walk is

$$\theta(T) = \frac{1}{2q(q+1)^2} (T_{s_1} + T_{s_1 s_0} + T_{s_0 s_1} + T_{s_0 s_1 s_0} + T_{s_2} + T_{s_2 s_0} + T_{s_0 s_2} + T_{s_0 s_2 s_0}).$$

By Proposition 5.7 we have $p^{(n)}(A, A) = \bar{p}^{(n)}(a, a)/(q+1)$. An asymptotic formula for $\bar{p}^{(n)}(a, a)$, and hence $p^{(n)}(A, A)$, can be computed using the techniques in [25]. The details of the calculation are somewhat involved, using the representation theory of the associated affine Hecke algebra. Thus we content ourselves here to simply stating the final result.

Theorem 5.8. *For the simple random walk on the set X_0 of cotype 0 simplices of an \tilde{A}_2 building we have*

$$p^{(n)}(A, A) \sim \frac{\sqrt{3}(q^2 + 4q - 1)^4}{\pi q(q+1)(q-1)^6} \left(\frac{q^2 + 4q - 1}{2q(q+1)} \right)^n n^{-4} \quad \text{as } n \rightarrow \infty$$

for all $A \in X_0$, where \sim denotes asymptotic equivalence.

References

- [1] P. Abramenko, K. Brown, *Transitivity properties for group actions on buildings*, J. Group Theory **10** (2007), 267–277.
- [2] P. Abramenko, K. Brown, *Buildings: Theory and Applications*, Graduate Texts in Mathematics **248**, Springer (2008).
- [3] P. Abramenko, J. Parkinson, H. Van Maldeghem, *A classification of commutative parabolic Hecke algebras*, J. Algebra **385** (2013), 115–133.
- [4] N.L. Biggs, A.G. Boshier, J. Shawe-Taylor, *Cubic distance-regular graphs*, J. London Math. Soc. (2) **33** (1986), 385–394.

- [5] W. Bosma, J. Cannon, C. Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), 235–265.
- [6] N. Bourbaki, *Lie Groups and Lie Algebras, Chapters 4–6*, Elements of Mathematics, Springer-Verlag, Berlin Heidelberg New York, (2002).
- [7] A. E. Brouwer, A.E. Cohen, *Computation of some parameters of Lie Geometries*, Ann. Discrete Math.: Algorithms in combinatorial design theory, North-Holland Math. Stud. **114** (1985), 1–48.
- [8] A. E. Brouwer, A. M. Cohen, A. Neumaier, *Distance Regular Graphs*, Springer-Verlag, Berlin, New York (1989).
- [9] P.-E. Caprace, C. Ciobotaru, *Gelfand pairs and strong transitivity for Euclidean buildings*, Ergod. Th. & Dynam. Sys. **35** (2015), 1056–1078.
- [10] D. I. Cartwright, *Harmonic functions on buildings of type \tilde{A}_n* , Random walks and discrete potential theory, M. Picardello and W. Woess (eds.), Symposia Mathematica **39**, Cambridge University Press, Cambridge, UK (1999), 104–138.
- [11] C. W. Curtis, N. Iwahori, R. Kilmoyer, *Hecke algebras and characters of parabolic type of finite groups with (B, N) -pairs*, Inst. Hautes Études Sci. Publ. Math. **40** (1971), 81–116.
- [12] S. Gaussent, P. Littelmann, *LS galleries, the path model, and MV cycles*, Duke Math. J. **127** (2005), 35–88.
- [13] Y. Gomi, *Structure constants of the Hecke algebras $\mathcal{H}(S, J)$ and applications*, J. Algebra **183** (1996), 514–544.
- [14] A.A. Ivanov, *Bounding the diameter of a distance-regular graph*, Sov. Math. Dokl. **28** (1983), 149–152.
- [15] N. Iwahori, *On the structure of the Hecke ring of a Chevalley group over a finite field*, J. Faculty Science, Tokyo University, **10** (1964), 215–236.
- [16] G. Lusztig, *Hecke algebras with unequal parameters*, CRM monograph series **18**, American Mathematical Society, 2003.
- [17] I. G. Macdonald, *Spherical functions on a group of p -adic type*, Ramanujan Institute for Advanced Mathematics, University of Madras, 1971.
- [18] I. G. Macdonald, *The Poincaré series of a Coxeter group*, Math. Ann. **199** (1972), 161–174.
- [19] H. Matsumoto, *Générateurs et relations des groupes de Weyl généralisés*, C. R. Acad. Sci. Paris **258** (1964), 3419–3422.
- [20] F. Miller Malley, *The Hall polynomial revisited*, J. Algebra **184** (1996), 363–371.
- [21] J. Parkinson, *Buildings and Hecke algebras*, J. Algebra **297** (2006), 1–49.
- [22] J. Parkinson, *Spherical harmonic analysis on affine buildings*, Math. Z. **253** (2006), 571–606.
- [23] J. Parkinson, *Isotropic random walks on affine buildings*, Ann. Inst. Fourier **57** (2007), 379–419.
- [24] J. Parkinson, A. Ram, C. Schwer, *Combinatorics in affine flag varieties*, J. Algebra **321** (2009), 3469–3493.
- [25] J. Parkinson, B. Schapira, *A local limit theorem for random walks on the chambers of \tilde{A}_2 buildings*, Progress in Probability **64**, Birkhäuser (2011), 15–53.
- [26] A. Ram, *Alcove walks, Hecke algebras, spherical functions, crystals and column strict tableaux*, Pure Appl. Math. Q. **2** (2006), 963–1013.
- [27] A. Schwartz, T. R. Shemanske, *Maximal orders in central simple algebras and Bruhat-Tits buildings*, J. Number Theory **56** (1996), 115–138.
- [28] C. Schwer, *Galleries, Hall-Littlewood polynomials, and structure constants of the spherical Hecke algebra*, Int. Math. Res. Not. Art. ID75395, (2006), 31pp.
- [29] A. Setyadi, *Distance in the affine buildings of SL_n and Sp_n* , Integers **8** (2008), A48, 17pp.
- [30] J. Tits, *Buildings of spherical type and finite BN -pairs*, Lecture Notes in Mathematics **386**, Springer-Verlag, Berlin-New York, 1974.

- [31] J. Tits, *Immeubles de type affine*, Buildings and the geometry of diagrams (Como, 1984), 159–190, Lecture Notes in Mathematics **1181**, Springer, Berlin, 1986.
- [32] R. Weiss, *The structure of affine buildings*, Annals of Mathematics Studies **168**, Princeton University Press, Princeton, NJ, 2009.
- [33] P.-H. Zieschang, *An algebraic approach to association schemes*, Lecture Notes in Math., **1628**, Springer, 1996.

Peter Abramenko
Department of Mathematics
University of Virginia
Charlottesville, VA 22904-4137, USA
pa8e@virginia.edu

James Parkinson
School of Mathematics and Statistics
University of Sydney
NSW, 2006, Australia
jamesp@maths.usyd.edu.au

Hendrik Van Maldeghem
Department of Mathematics
Ghent University
Krijgslaan 281, S22,
9000 Ghent, Belgium
hvm@cage.UGent.be